

ON BOUNDARIES OF COXETER GROUPS AND TOPOLOGICAL FRACTAL STRUCTURES

TETSUYA HOSAKA

ABSTRACT. In this paper, based on research on rank-one isometries by W. Ballmann and M. Brin and recent research on rank-one isometries of Coxeter groups by P. Caprace and K. Fujiwara, we study a topological fractal structure of boundaries of Coxeter groups. We also show that the limit-point set is dense in a boundary of a Coxeter group and introduce some observations on boundaries of CAT(0) groups with rank-one isometries.

1. INTRODUCTION

In this paper, we study boundaries of Coxeter groups, where we suppose that Coxeter groups are finitely generated and infinite. A Coxeter group acts geometrically (i.e. properly and cocompactly by isometries) on a Davis complex which is a CAT(0) space [28] and every Coxeter group is a CAT(0) group. Details of Coxeter groups and Coxeter systems are found in [5], [7], [13], [23] and [31], and details of CAT(0) spaces, CAT(0) groups and their boundaries are found in [6], [9] and [16].

Now we suppose that an infinite group G acts geometrically on a proper CAT(0) space X and G is non-elementary (hence $|\partial X| > 2$).

A hyperbolic isometry g of a proper CAT(0) space X is said to be *rank-one*, if some (any) axis for g does not bound a flat half-plane. In [1, Theorem A], W. Ballmann and M. Brin have proved that if there exists a rank-one isometry $g \in G$ of X then for any two non-empty open subsets U and V of ∂X , there exists an element $g \in G$ such that $g(\partial X - U) \subset V$ and $g^{-1}(\partial X - V) \subset U$ where it is possible to choose g to be rank-one (cf. [8], [18]).

Date: November 29, 2010.

2000 Mathematics Subject Classification. 20F65; 20F55; 57M07.

Key words and phrases. Coxeter group; boundary; CAT(0) space; Davis complex; rank-one isometry; minimal; fractal.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 21740037).

This statement implies that if there exists a rank-one isometry $g \in G$ of X then we can say that the boundary ∂X has a *topological fractal structure*; that is, for any proper closed subset F of ∂X and any non-empty open subset U of ∂X , there exists $g \in G$ such that $gF \subset U$.

We first note that if G is hyperbolic then G contains a rank-one isometry and the boundary ∂X has a topological fractal structure.

In particular, if G is hyperbolic and the boundary ∂X is an n -sphere then the boundary $\partial X \approx \mathbb{S}^n$ has a topological fractal structure. This case is the most simple case of boundaries of CAT(0) groups with rank-one isometries. In general, the boundary ∂X with a topological fractal structure is very complex.

In [15], H. Fischer has investigated the boundary $\partial \Sigma$ of the Davis complex of a right-angled Coxeter group whose nerve is a connected closed orientable PL-manifold. These boundaries are typical examples of boundaries with topological fractal structures. If the boundary ∂X with a topological fractal structure contains some proper closed subset F which has a something non-trivial homotopy type, then any (small) open subset U of ∂X contains gF for some homeomorphism $g \in G$ of ∂X and $\{gF \mid g \in G\}$ is dense in ∂X , where every gF is homeomorphic to F .

Also for a proper closed subset F of the boundary ∂X with a topological fractal structure such that the complement $\partial X - F$ is a very small neighborhood, any (small) open subset U of ∂X contains gF for some homeomorphism $g \in G$ of ∂X .

Thus, in such a case that G contains a rank-one isometry and ∂X is not an n -sphere, then the boundary ∂X seems to be just a topological fractal.

This fractal structure seems to be suggested in some research on boundaries of CAT(0) groups by M. Bestvina (cf. [4]) and some research on cohomology of boundaries of Coxeter groups (cf. [3], [11], [14], [19]).

If the boundary ∂X has a topological fractal structure, then (the action of G on) ∂X is *minimal*; that is, every orbit $G\alpha$ is dense in the boundary ∂X . Indeed if we take $F = \{\alpha\}$ then for any open subset U of ∂X , $gF \subset U$ for some $g \in G$.

Also then (the action of G on) ∂X is *scrambled*; that is, for any two points $\alpha, \beta \in \partial X$ with $\alpha \neq \beta$,

$$\begin{aligned} \limsup \{d_{\partial X}(g\alpha, g\beta) \mid g \in G\} &> 0 \text{ and} \\ \liminf \{d_{\partial X}(g\alpha, g\beta) \mid g \in G\} &= 0 \end{aligned}$$

(cf. [21]). Indeed $\limsup \{d_{\partial X}(g\alpha, g\beta) \mid g \in G\} > 0$ always holds ([21, Theorem 3.1]) and if we take $F = \{\alpha, \beta\}$ then for any small open subset

U of ∂X , $gF \subset U$ for some $g \in G$, hence $\liminf\{d_{\partial X}(g\alpha, g\beta) \mid g \in G\} = 0$.

Thus if the boundary ∂X is a topological fractal, then ∂X is minimal and scrambled.

We can find recent research on minimality and scrambled sets of boundaries of Coxeter groups in [20] and [21].

From recent research on rank-one isometries of Coxeter groups by P. Caprace and K. Fujiwara [8, Proposition 4.5], we obtain that for a Coxeter system (W, S) such that S is finite and W is infinite and non-elementary, if (W, S) is irreducible and non-affine then the Coxeter group W contains a rank-one isometry of the Davis complex Σ defined by (W, S) . Hence a finitely generated, infinite and non-elementary Coxeter group W contains a rank-one isometry if and only if W does not contain a finite-index subgroup which splits as a product $W_1 \times W_2$ where W_1 and W_2 are infinite.

By the observation above, we obtain the following theorem.

Theorem 1.1. *Let (W, S) be a Coxeter system such that W is infinite and non-elementary and S is finite. For the Davis complex Σ of (W, S) and any proper $CAT(0)$ space X on which W acts geometrically, the following statements are equivalent.*

- (1) $(W_{\tilde{S}}, \tilde{S})$ is irreducible and non-affine.
- (2) W contains a rank-one isometry of Σ .
- (3) W contains a rank-one isometry of X .
- (4) $\partial\Sigma$ has a topological fractal structure.
- (5) $\partial\Sigma$ is minimal.
- (6) $\partial\Sigma$ is scrambled.
- (7) ∂X has a topological fractal structure.
- (8) ∂X is minimal.
- (9) ∂X is scrambled.
- (10) Σ does not contain a quasi-dense subspace which splits as a product $\Sigma_1 \times \Sigma_2$ of two unbounded subspaces.
- (11) X does not contain a quasi-dense subspace which splits as a product $X_1 \times X_2$ of two unbounded subspaces.
- (12) W does not contain a finite-index subgroup which splits as a product $W_1 \times W_2$ of two infinite subgroups.

Here $W_{\tilde{S}}$ is the minimum finite-index parabolic subgroup of (W, S) ([13], cf. [20], [21]).

Thus if (W, S) is an irreducible Coxeter system, then W is finite, W is affine or W contains a rank-one isometry.

Hence for any Coxeter system (W, S) and the irreducible decomposition of (W, S) as

$$W = W_{S_1} \times \cdots \times W_{S_k} \times W_{S_{k+1}} \times \cdots \times W_{S_n},$$

each W_{S_i} is finite, affine or contains a rank-one isometry.

It is known that the following problem is open.

Question. Suppose that a group G acts geometrically on a proper CAT(0) space X . Then is it the case that the limit-point set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is dense in the boundary ∂X ?

Here g^∞ is the limit-point of the boundary ∂X to which the sequence $\{g^i x_0 \mid i \in \mathbb{N}\} \subset X$ converges in $X \cup \partial X$, where x_0 is a point of X and the limit-point g^∞ is not depend on x_0 . We note that any element g of a CAT(0) group G with the order $o(g) = \infty$ is a hyperbolic isometry.

We obtain a positive answer to this question for Coxeter groups.

Theorem 1.2. *Suppose that a finitely generated infinite Coxeter group W acts geometrically on a proper CAT(0) space X . Then the limit-point set $\{w^\infty \mid w \in W, o(w) = \infty\}$ is dense in the boundary ∂X .*

Finally, we introduce some observations on boundaries of CAT(0) groups with rank-one isometries in Section 4, which relates to local properties of boundaries of CAT(0) groups.

2. RANK-ONE ISOMETRIES OF COXETER GROUPS AND TOPOLOGICAL FRACTAL STRUCTURES OF THEIR BOUNDARIES

We prove Theorem 1.1.

Proof of Theorem 1.1. We first obtain the equivalence (1) $\Leftrightarrow$ (2) $\Leftrightarrow$ (12) from [8, Proposition 4.5] and the observation in Section 1. Also (2) $\Leftrightarrow$ (3) holds by [1, Theorem B].

From the observation in Section 1 on rank-one isometries and topological fractal structures of boundaries, we obtain (2) $\Rightarrow$ (4), (4) $\Rightarrow$ (5) and (4) $\Rightarrow$ (6), also, (3) $\Rightarrow$ (7), (7) $\Rightarrow$ (8) and (7) $\Rightarrow$ (9).

Concerning scrambled sets of boundaries, [21, Theorem 5.5] implies (6) $\Rightarrow$ (10) and (9) $\Rightarrow$ (11).

Also concerning minimality of boundaries, [20, Theorem 6.4] implies (5) $\Rightarrow$ (12) and (8) $\Rightarrow$ (12).

By splitting theorems (cf. [22], [27]), we obtain (10) $\Rightarrow$ (12) and (11) $\Rightarrow$ (12) (cf. [20, Proposition 6.3]).

Therefore the statements (1)–(12) are equivalent. $\square$

3. ON LIMIT-POINT SETS OF BOUNDARIES OF COXETER GROUPS

We prove Theorem 1.2.

Proof of Theorem 1.2. Suppose that a finitely generated infinite Coxeter group W acts geometrically on a proper CAT(0) space X .

Here there exists $S \subset W$ such that (W, S) is a Coxeter system. Now we consider the irreducible decomposition of (W, S) as

$$W = W_{S_1} \times \cdots \times W_{S_k} \times W_{S_{k+1}} \times \cdots \times W_{S_n}$$

where each (W_i, S_i) is irreducible and we may suppose that W_{S_i} is infinite for any $i = 1, \dots, k$ and W_{S_i} is finite for any $i = k+1, \dots, n$. Let $W' = W_{S_1} \times \cdots \times W_{S_k}$. Then W' is a finite-index subgroup of W and acts geometrically on the CAT(0) space X (where W' is the minimum finite-index parabolic subgroup of (W, S)).

Here we note that every Coxeter group has finite center. Hence by the splitting theorem [22, Theorem 2] and [27, Corollary 10], X contains a closed convex W' -invariant quasi-dense subspace X' which splits as a product $X' = X_1 \times \cdots \times X_k$ where the action of $W' = W_{S_1} \times \cdots \times W_{S_k}$ on $X' = X_1 \times \cdots \times X_k$ splits and W_{S_i} acts geometrically on X_i for each $i = 1, \dots, k$.

Then every irreducible infinite Coxeter group W_{S_i} is either affine or contains a rank-one isometry by [8, Proposition 6.5] and the observation in Section 1.

If W_{S_i} is affine, then W_{S_i} contains a finite-index subgroup which is isomorphic to $\mathbb{Z}^{n_i}$ and X_i contains a quasi-dense subspace which is isometric to $\mathbb{R}^{n_i}$. Hence the limit-point set $\{w_i^\infty \mid w_i \in W_i, o(w_i) = \infty\}$ is dense in the boundary ∂X_i .

Also if W_{S_i} contains a rank-one isometry, then the action of W_{S_i} on the boundary ∂X_i is minimal. Hence [20, Proposition 6.2] implies that the limit-point set $\{w_i^\infty \mid w_i \in W_i, o(w_i) = \infty\}$ is dense in the boundary ∂X_i .

Therefore, by a similar argument to the proof of [20, Proposition 6.5], we obtain that the limit-point set $\{w^\infty \mid w \in W, o(w) = \infty\}$ is dense in the boundary ∂X . $\square$

4. OBSERVATIONS ON BOUNDARIES OF CAT(0) GROUPS WITH RANK-ONE ISOMETRIES

We introduce some observations on boundaries of CAT(0) groups with rank-one isometries.

Now we suppose that a group G acts geometrically on a proper CAT(0) space X and suppose that G contains a rank-one isometry (hence the boundary ∂X has a topological fractal structure).

Let V be a non-empty open subset of ∂X whose closure $\text{cl} V$ is a proper subset of ∂X . Then there exists a rank-one isometry $g \in G$ as $g^\infty \in V$, because the limit-point set of rank-one isometries in G is dense in ∂X . Indeed ∂X is minimal and

$$Gg^\infty = \{ag^\infty \mid a \in G\} = \{(aga^{-1})^\infty \mid a \in G\}$$

is dense in the boundary ∂X .

Every rank-one isometry acts with *north-south dynamics* on the boundary ∂X (cf. [18, p.7]). Hence, since g is a rank-one isometry of X and $g^\infty \in V$, the set $\{g^i V \mid i \in \mathbb{N}\}$ is a neighborhood basis for g^∞ in ∂X . Here all $g^i V$ are homeomorphic to V .

Thus if there exists a non-empty open subset V of ∂X whose closure $\text{cl} V$ is a proper subset of ∂X such that V has some *topological property* (P) , then ∂X has the locally *topological property* (P) at the limit-point g^∞ .

Also for any rank-one isometry $h \in G$, we can consider the limit-point $h^\infty \in \partial X$. Then Gh^∞ is dense in ∂X , since ∂X is minimal. Hence $ah^\infty \in V$ for some $a \in G$. Then $h^\infty \in a^{-1}V$ and $a^{-1}V$ is homeomorphic to V . Thus the boundary ∂X has the locally *topological property* (P) at the limit-point h^∞ of all rank-one isometries $h \in G$.

We also note that the limit-point set of all rank-one isometries is dense in the boundary ∂X .

As one example, if there exists a non-empty *connected* open subset V of ∂X whose closure $\text{cl} V$ is a proper subset of ∂X , then ∂X is locally *connected* at the limit-points g^∞ of all rank-one isometries $g \in G$.

Moreover if ∂X is non-locally connected at some point $\alpha \in \partial X$, then ∂X is non-locally connected at $g\alpha$ for all $g \in G$. Here $G\alpha$ is also dense in ∂X .

It seems that these arguments relate to research on local connectivity of boundaries of CAT(0) groups by M. Mihalik, K. Ruane and S. Tschantz ([25], [26]) and research on cut-points and limit-points of boundaries of CAT(0) groups by P. Papasoglu and E. L. Swenson ([29], [30]).

Also as one application, we obtain the following theorem by a similar argument to the proof of [24, Theorem 4.4].

Theorem 4.1. *If a CAT(0) group G with a rank-one isometry acts geometrically on a proper CAT(0) space X , then the following statements are equivalent:*

- (i) *the boundary ∂X is an n -manifold,*
- (ii) *the boundary ∂X of X contains some closed neighborhood U which is homeomorphic to an n -ball,*

(iii) *the boundary ∂X is homeomorphic to an n -sphere.*

Proof. We first note that the implications (iii) $\Rightarrow$ (i) $\Rightarrow$ (ii) are obvious.

Hence now we show the implication (ii) $\Rightarrow$ (iii).

Suppose that (iii) holds; that is, the boundary ∂X of X contains some closed neighborhood U which is homeomorphic to an n -ball. For a point $\alpha \in \partial X - U$, there exists $g \in G$ such that $g\alpha \in \text{Int } U$, since the action of G on ∂X is minimal. Then $V := g^{-1}U$ is a neighborhood of α which is homeomorphic to an n -ball. Let U' and V' be proper subsets of $\text{Int } U$ and $\text{Int } V$ respectively such that U' and V' are homeomorphic to an n -ball and $U' \cap V' = \emptyset$. Let $F = \partial X - \text{Int } U'$. Then there exists $g' \in G$ such that $g'F \subset V'$, because the boundary ∂X has a topological fractal structure. Then $g'U' \cup V' = \partial X$ and $g'U'$ and V' are homeomorphic to an n -ball. (Moreover, $g'U \cup V = \partial X$ and $g'U$ and V are homeomorphic to an n -ball.) Using some argument on bicollars of n -disks, we obtain that ∂X is homeomorphic to an n -sphere. $\square$

Acknowledgement. The author would like to thank Dr. Naotsugu Chinen for helpful discussion and helpful advice.

REFERENCES

- [1] W. Ballmann and M. Brin, *Orbihedra of nonpositive curvature*, Inst. Hautes Études Sci. Publ. Math. 82 (1995), 169–209.
- [2] W. Ballmann, M. Gromov and V. Schroeder, *Manifolds of Nonpositive Curvature*, Progr. Math. vol. 61, Birkhäuser, Boston MA, 1985.
- [3] M. Bestvina, *The virtual cohomological dimension of Coxeter groups*, Geometric Group Theory Vol. 1, LMS Lecture Notes, vol. 181, 1993, pp. 19–23.
- [4] M. Bestvina, *Local homology properties of boundaries of groups*, Michigan Math. J. 43 (1996), 123–139.
- [5] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV–VI, Masson, Paris, 1981.
- [6] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [7] K. S. Brown, *Buildings*, Springer-Verlag, 1980.
- [8] P. Caprace and K. Fujiwara, *Rank-one isometries of buildings and quasi-morphisms of Kac-Moody groups*, Geom. Funct. Anal. 19 (2010), 1296–1319.
- [9] C. B. Croke and B. Kleiner, *Spaces with nonpositive curvature and their ideal boundaries*, Topology 39 (2000), 549–556.
- [10] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. 117 (1983), 293–324.
- [11] M. W. Davis, *The cohomology of a Coxeter group with group ring coefficients*, Duke Math. J. 91 (no.2) (1998), 297–314.
- [12] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.

- [13] V. Deodhar, *On the root system of a Coxeter group*, Commun. Algebra 10 (1982), 611–630.
- [14] A. N. Dranishnikov, *On the virtual cohomological dimensions of Coxeter groups*, Proc. Amer. Math. Soc. 125 (no.7) (1997), 1885–1891.
- [15] H. Fischer, *Boundaries of right-angled Coxeter groups with manifold nerves* Topology 42 (2003), 423–446.
- [16] E. Ghys and P. de la Harpe (ed), *Sur les Groupes Hyperboliques d’après Mikhael Gromov*, Progr. Math. vol. 83, Birkhäuser, Boston MA, 1990.
- [17] M. Gromov, *Hyperbolic groups*, in Essays in group theory (Edited by S. M. Gersten), pp. 75–263, M.S.R.I. Publ. 8, 1987.
- [18] U. Hamenstädt, *Rank-one isometries of proper $CAT(0)$ -spaces*, Contemp. Math. 501 (2009), 43–59.
- [19] T. Hosaka, *On the cohomology of Coxeter groups*, J. Pure Appl. Algebra 162 (2001), 291–301.
- [20] T. Hosaka, *Minimality of the boundary of a right-angled Coxeter system*, Proc. Amer. Math. Soc., 137 (2009), 899–910.
- [21] T. Hosaka, *$CAT(0)$ groups and Coxeter groups whose boundaries are scrambled sets*, J. Pure Appl. Algebra 214 (2010), 919–936.
- [22] T. Hosaka, *On splitting theorems for $CAT(0)$ spaces and compact geodesic spaces of non-positive curvature*, arXiv:math.GR/0405551, preprint.
- [23] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [24] I. Kapovich and N. Benakli, *Boundaries of hyperbolic groups*, in Combinatorial and Geometric Group Theory (R. Gilman et al, editors), Contemporary Mathematics vol. 296, pp. 39–94, 2002.
- [25] M. Mihalik and K. Ruane, *$CAT(0)$ groups with non-locally connected boundary*, J. London Math. Soc. (2) 60 (1999), 757–770.
- [26] M. Mihalik, K. Ruane and S. Tschant, *Local connectivity of right-angled Coxeter group boundaries*, J. Group Theory 10 (2007), 531–560.
- [27] N. Monod, *Superrigidity for irreducible lattices and geometric splitting*, J. Amer. Math. Soc. 19 (2006), 781–814.
- [28] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, Ohio State University, 1988.
- [29] P. Papasoglu and E. L. Swenson, *Boundaries and JSJ decompositions of $CAT(0)$ -groups*, Geom. Funct. Anal. 19 (2009), 558–590.
- [30] E. L. Swenson, *A cut point theorem for $CAT(0)$ groups*, J. Differential Geom. 53 (1999), 327–358.
- [31] J. Tits, *Le problème des mots dans les groupes de Coxeter*, Symposia Mathematica, vol. 1, pp. 175–185, Academic Press, London, 1969.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, UTSUNOMIYA
UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN
E-mail address: hosaka@cc.utsunomiya-u.ac.jp